

ADDITION AND SUBSPACE THEOREMS FOR ASYMPTOTIC LARGE INDUCTIVE DIMENSION

T.RADUL

ABSTRACT. We prove addition and subspace theorems for asymptotic large inductive dimension. We investigate a transfinite extension of this dimension and show that it is trivial.

0. Asymptotic dimension asdim of a metric space was defined by Gromov for studying asymptotic invariants of discrete groups [1]. This dimension can be considered as asymptotic analogue of the Lebesgue covering dimension dim . Dranishnikov has introduced dimension asInd which is analogous to large inductive dimension Ind [2].

Some of basic theorems of classical dimension theory are sum, addition and subspace theorems for different dimensions and classes of topological spaces. Here we mention some of them related to dimension Ind . (All mentioned facts from classic dimension theory could be found in [3]).

1. Countable sum theorem: If a strongly hereditarily normal space X can be represented as the union of a sequence $F_1, F_2, \dots$ of closed subsets such that $\text{Ind } F_i \leq n$ for each $i \in \mathbb{N}$, then $\text{Ind } X \leq n$.

2. Addition theorem: If a hereditarily normal space X is represented as the union of two subspaces X_1 and X_2 , then $\text{Ind } X \leq \text{Ind } X_1 + \text{Ind } X_2 + 1$.

Let us remark that we cannot generalize the above theorems for the class of all normal spaces. We have only a weaker result:

3. If a normal space X is represented as the union of two closed subsets X_1 and X_2 , then $\text{Ind } X \leq \text{Ind } X_1 + \text{Ind } X_2$.

And finally:

1991 *Mathematics Subject Classification.* 54F45, 54D35.

Key words and phrases. Asymptotic dimension, transfinite extension.

4. Subspace theorem: For each closed subset M of a normal space X we have $\text{Ind } M \leq \text{Ind } X$. Let us remark that if X is a strongly hereditarily normal space, then the condition that M is closed could be dropped.

There is no countable sum theorem for asInd . Indeed, the space of integers $\mathbb{Z}$ has asymptotic dimension 1 but it is countable union of its points which have asymptotic dimension -1 . However we prove subspace and addition theorem for asInd in this paper.

Extending codomain of Ind to ordinal numbers we obtain the transfinite extension trInd of the dimension Ind . It is known that there exists a space S_α such that $\text{trInd } S_\alpha = \alpha$ for each countable ordinal number α . Zarichnyi has proposed to consider transfinite extension of asInd and conjectured that this extension is trivial. We prove this conjecture: if a space has a transfinite asymptotic dimension, then its dimension is finite.

The paper is organized as follows: in Section 1 we give some necessary definitions and introduce some denotations, in Section 2 we prove some theorem which can be consider as a weak version of countable sum theorem, in Section 3 we obtain the main results and in Section 4 we show that the transfinite extension of asInd is trivial.

1. Let $A_1, A_2 \subset X$ be two disjoint closed subsets in a topological space. We recall that a *partition* between A_1 and A_2 is a subset $C \subset X$ such that there are open disjoint sets U_1, U_2 satisfying the conditions: $X \setminus C = U_1 \cup U_2$, $A_1 \subset U_1$ and $A_2 \subset U_2$. Clearly a partition C is a closed subset of X .

We recall the definition of the large inductive dimension Ind [3]: $\text{Ind } X = -1$ iff $X = \emptyset$; $\text{Ind } X \leq n$ if for every two disjoint closed subsets $A_1, A_2 \subset X$ there is a partition C with $\text{Ind } C \leq n - 1$.

We will define the dimension asInd for the class of proper metric space. We recall that a metric space is *proper* if every closed ball is compact. We assume that there is chosen some base point $x_0 \in X$ for each proper metric space X . The generic metric we denote by d . If X is a metric space and $A \subset X$ we denote by $N_r(A)$ the closed r -neighborhood of A in X : $N_r(A) = \{x \in X \mid d(x, A) \leq r\}$ and by $B_r(A)$ the open r -neighborhood: $B_r(A) = \{x \in X \mid d(x, A) < r\}$.

A subset W of a metric space is called an *asymptotic neighborhood* of a set $A \subset X$ if $\lim_{r \rightarrow \infty} d(A, B_r(x_0) \setminus W) = \infty$. We call two subsets $A, A' \subset X$ in a

metric space X *asymptotically disjoint* if $\lim_{r \rightarrow \infty} d(A_1 \setminus B_r(x_0), A_2 \setminus B_r(x_0)) = \infty$.

It is easy to see that sets A_1 and A_2 are asymptotically disjoint iff $X \setminus A_2$ is an asymptotic neighborhood of A_1 and $X \setminus A_1$ is an asymptotic neighborhood of A_2 .

A map $\phi : X \rightarrow I = [0, 1]$ is called *slowly oscillating* if for any $r > 0$, for given $\varepsilon > 0$ there exists $D > 0$ such that $\text{diam } \phi(B_r(x)) < \varepsilon$ for any x with $d(x, x_0) \geq D$. If $C_h(X)$ is the set of all continuous slow oscillating functions $\phi : X \rightarrow I$, then the *Higson compactification* is the closure of the image of X under the embedding $\Phi : X \rightarrow I^{C_h(X)}$ defined as $\Phi(x) = (\phi(x) \mid \phi \in C_h(X)) \in I^{C_h(X)}$. We denote the Higson compactification of a proper metric space X by cX and the remainder $cX \setminus X$ by νX . The compactum νX is called *Higson corona*. Let us remark that νX does not need to be metrizable.

Let C be a subset of a proper metric space X . By C' we denote the intersection $\text{cl } C \cap \nu X$ of the closure $\text{cl } C$ in the Higson compactification cX . Clearly, two sets A_1 and A_2 are asymptotically disjoint iff their traces A'_1 and A'_2 in the Higson corona are disjoint. Note that for each $r > 0$ we have $N_r(C)' = C'$.

Let $A_1, A_2 \subset X$ be two asymptotically disjoint subsets of a proper metric space X . A closed subset $C \subset X$ is called an *asymptotic separator* for A_1 and A_2 if its trace C' is a partition for A'_1 and A'_2 in νX .

We define $\text{asInd } X = -1$ if and only if X is bounded; $\text{asInd } X \leq n$ if for every two asymptotically disjoint sets $A, B \subset X$ there is an asymptotic separator C with $\text{asInd } C \leq n - 1$. Naturally we say $\text{asInd } X = n$ if $\text{asInd } X \leq n$ and it is not true that $\text{asInd } X \leq n - 1$. We set $\text{asInd } X = \infty$ if $\text{asInd } X > n$ for each $n \in \mathbb{N}$ [2].

For each $A \subset Y \subset X$ we denote by $\text{ex}_Y A = Y' \setminus (Y \setminus A)'$. Clearly, $\text{ex}_Y A$ is an open set in Y' . By ω^ω we denote the set of all functions $\tau : \mathbb{N} \cup \{0\} \rightarrow \mathbb{N} \cup \{0\}$ such that $\tau(0) = 0$. For each $\tau \in \omega^\omega$ we define the set $V_\tau^Y(A)$ as follows: $V_\tau^Y(A) = \{y \in Y \mid d(y, x_0) \geq \tau([d(y, A)])\}$ where $[d(y, A)]$ is the enter part of real number $d(y, A)$. Clearly, $A \subset V_\tau^Y(A)$ for each $\tau \in \omega^\omega$. If $X = Y$ we use simpler denotations $\text{ex } A$ and $V_\tau(A)$.

Lemma 1. *The family $\{\text{ex}(V_\tau(A)) \mid \tau \in \omega^\omega\}$ forms the base of neighborhoods of the set A' in the space νX .*

Proof. Let us show firstly that $A' \subset \text{ex}(V_\tau(A))$ for each $\tau \in \omega^\omega$. It is enough to show that A and $X \setminus V_\tau(A)$ are asymptotically disjoint. Fix any $D > 0$. Put

$R = \max\{\tau(l) \mid l \in \{0, \dots, [D]\}\}$. Then for each $x \in (X \setminus V_\tau(A)) \setminus B_R(x_0)$ we have $R \leq d(x, x_0) < \tau([d(x, A)])$. Then $d(x, A) \geq [D] + 1 > D$.

Consider now any closed subset B of νX such that $B \cap A' = \emptyset$. Then there exists a continuous function $f : \nu X \rightarrow [0, 1]$ such that $f(B) \subset \{0\}$ and $f(A') \subset \{1\}$. We can extend f to a continuous function $g : cX \rightarrow [0, 1]$ such that $A \subset g^{-1}(1)$. Put $C = g^{-1}[0, \frac{1}{2}] \cap X$. The sets C and A are asymptotically disjoint. For each $n \in \mathbb{N}$ there exists $R(n) > 0$ such that $d(C \setminus B_{R(n)}(x_0), A) \geq n$. Put $\tau(0) = 0$ and $\tau(n) = [R(n+1)]$. Choose any $c \in C$. Then $d(c, x_0) < R([d(c, A)] + 1) = \tau([d(c, A)])$. So, $C \subset X \setminus V_\tau(A)$ and $B \subset C' \subset (X \setminus V_\tau(A))' \subset \nu X \setminus \text{ex } V_\tau(A)$. Hence $\{\text{ex}(V_\tau(A)) \mid \tau \in \omega^\omega\}$ forms the base of neighborhoods of the set A' in the space νX and the lemma is proved.

2. Let X be a proper metric space and X_0 is an unbounded subset of X . We say that X_0 is a *kernel* of X if there exists a sequence $(k_i)_{i=0}^\infty$ of natural numbers such that $k_i \rightarrow \infty$ and for each $i \in \mathbb{N} \cup \{0\}$, $x \in X \setminus N_i(X_0)$ we have $B_{k_i}(x) = \{x\}$. We suppose that $x_0 \in X_0$ where x_0 is the base point in X .

Lemma 2. *If X_0 is a kernel of a metric proper space X then the family $\{V_\tau(X_0)\} \mid \tau \in \omega^\omega\}$ forms the base of clopen neighborhoods of the set X'_0 in the space νX .*

Proof. It follows from Lemma 1 that it is enough to prove that $V_\tau(X_0)' \cap (X \setminus V_\tau(X_0))' = \emptyset$. Suppose the contrary: there exist $x \in V_\tau(X_0)' \cap (X \setminus V_\tau(X_0))'$. Let U be a neighborhood of x in cX . Then there exist two sequences (a_i) and (b_i) in $V_\tau(X_0) \cap U$ and $(X \setminus V_\tau(X_0)) \cap U$ respectively such that $0 < d(a_i, b_i) \leq r$ for some $r > 0$ and $a_i, b_i \in X \setminus B_i(x_0)$. Choose any $n_0 \in \mathbb{N}$ such that $k_n > r$ for each $n \geq n_0$ where (k_n) is a sequence from the definition of kernel. Then $a_i, b_i \in N_{n_0}(X_0)$ for each $i \in \mathbb{N}$. Hence $\emptyset \neq \text{cl } U \cap N_{n_0}(X_0)' = \text{cl } U \cap X'_0$ and $x \in X_0$. We obtain the contradiction.

Lemma 3. *If X_0 is a kernel of X then $\text{Ind}(\nu X \setminus V_\tau(X_0))' \leq 0$ for each $\tau \in \omega^\omega$.*

Proof. Since $\nu X \setminus V_\tau(X_0)'$ is compact, it is enough to prove that the space $\nu X \setminus V_\tau(X_0)'$ has a base of clopen sets [3, Theorem 1.6.5].

Choose any $x \in \nu X \setminus V_\tau(X_0)'$ and its open neighborhood $U \subset \nu X \setminus V_\tau(X_0)'$. Take a continuous function $f : \nu X \rightarrow [0, 1]$ such that $f(x) = 0$ and $f(\nu X \setminus U) \subset \{1\}$. Extend f to the continuous function $g : cX \rightarrow [0, 1]$. Put $A = g^{-1}[0, \frac{1}{2}] \cap X$.

and $C = g^{-1}[\frac{2}{3}, 1] \cap X$. The sets C and A are asymptotically disjoint. Moreover, $x \in A'$ and $\nu X \setminus U \subset C'$. For each $n \in \mathbb{N}$ there exists $R(n) > 0$ such that $d(C \setminus B_R(x_0), A \setminus B_R(x_0)) \geq n$. Put $\tau(0) = 0$ and $\tau(n) = [R(n+1)]$. We can show that $V_\tau(A)'$ is a clopen neighborhood of x such that $V_\tau(A)' \subset U$ using the same reasoning as in Lemmas 1 and 2. The lemma is proved.

Let us define a preorder $\leq^*$ in ω^ω as follows $\tau \leq^* \sigma$ iff there exists $n \in \mathbb{N}$ such that $\tau(i) \leq \sigma(i)$ for each $i \geq n$. It is easy to check that $V_\tau(A)' \subset V_\sigma(A)'$ if $\sigma \leq^* \tau$.

Theorem 1. *Let X_0 be a kernel of a metric proper space X such that $\text{asInd } X_0 \leq k \geq 0$. Then $\text{asInd } X \leq k$.*

Proof. We use induction respect to k . Let $\text{asInd } X_0 \leq 0$. Then $\text{Ind } X'_0 \leq 0$ [2]. Consider any two asymptotically disjoint sets A and B in X . We can represent $X'_0 = K \cup L$ where K and L are two disjoint closed subsets of X'_0 such that $A' \cap X'_0 \subset K$ and $B' \cap X'_0 \subset L$. Then $A' \cup K$ and $B' \cup L$ are two disjoint closed subsets of νX . Choose two disjoint open subsets U_1 and U_2 of νX such that $A' \cup K \subset U_1$ and $B' \cup L \subset U_2$. Then $U_1 \cup U_2$ is a neighborhood of X_0 in νX and there exists $\tau \in \omega^\omega$ such that $V_\tau(X'_0)' \subset U_1 \cup U_2$. The sets $A' \cap (\nu X \setminus V_\tau(X'_0)')$ and $B' \cap (\nu X \setminus V_\tau(X'_0)')$ are two disjoint closed subset of the 0-dimensional space $\nu X \setminus V_\tau(X'_0)'$, so, there exist two open disjoint subsets O_1, O_2 of $\nu X \setminus V_\tau(X'_0)'$ such that $A' \cap (\nu X \setminus V_\tau(X'_0)') \subset O_1$, $B' \cap (\nu X \setminus V_\tau(X'_0)') \subset O_2$ and $O_1 \cup O_2 = \nu X \setminus V_\tau(X'_0)'$. Put $V_1 = (U_1 \cap V_\tau(X'_0)') \cup O_1$ and $V_2 = (U_2 \cap V_\tau(X'_0)') \cup O_2$. Since $V_\tau(X'_0)'$ is a clopen subset of νX , the sets V_1 and V_2 are open. Moreover, the sets V_1 and V_2 are disjoint, $V_1 \cup V_2 = \nu X$ and $A' \subset V_1$ and $B' \subset V_2$. Thus, empty space is a partition between A' and B' in νX . Hence, empty space is an asymptotic separator between A and B in X and $\text{asInd } X \leq 0$.

Suppose that the theorem is proved for each $i < n \geq 1$. Consider the case when $\text{asInd } X_0 \leq n$. Let A and B be any asymptotically disjoint subset of X . Then $A' \cap X'_0$ and $B' \cap X'_0$ are disjoint closed subsets of X'_0 and we can choose a continuous function $f' : X'_0 \rightarrow [0, 1]$ such that $f'(A' \cap X'_0) \subset \{0\}$ and $f'(B' \cap X'_0) \subset \{1\}$. We can extend this function to a continuous function $f : \text{cl } X_0 \rightarrow [0, 1]$. The sets $A_1 = (f^{-1}[0, \frac{1}{3}]) \cap X_0$ and $B_1 = (f^{-1}[\frac{2}{3}, 1]) \cap X_0$ are asymptotically disjoint and we can choose an asymptotic separator $L \subset X_0$ between them such that $\text{asInd } L < n$.

Represent $X = \bigcup_{i=0}^{\infty} X_i$ where $X_i = N(X_0) \setminus N_{i-1}(X_0)$ for $i \in \mathbb{N}$. It follows from

the definition of kernel that for each $R > 0$ there exists $i(R) \in \mathbb{N}$ such that the set $\bigcup_{k=i(R)}^\infty X_k$ is R -discrete and $d(\bigcup_{k=i(R)}^\infty X_k, \bigcup_{k=0}^{i(R)-1} X_k) \geq R$.

Since L is an asymptotic separator in X_0 between A_1 and B_1 , then L' is a partition in X'_0 between A'_1 and B'_1 . Then we can choose two open disjoint sets O_A and O_B in X'_0 such that $A'_1 \subset O_A$, $B'_1 \subset O_B$ and $X'_0 \setminus L' = O_A \cup O_B$.

For each $\tau \in \omega^\omega$ we can represent $X_0 \setminus V_\tau(L)$ as a union of two disjoint sets A_τ and B_τ such that $A'_\tau \subset O_A$ and $B'_\tau \subset O_B$. Moreover, we can suppose that for each $\tau \leq^* \sigma$ there exists $R > 0$ such that $A_\sigma \supset A_\tau \setminus B_R(x_0)$ and $B_\sigma \supset B_\tau \setminus B_R(x_0)$. Define for each $\tau \in \omega^\omega$ two subsets C_τ, D_τ in X as follows: $C_\tau = \bigcup_{i=0}^\infty \{x \in X_i \mid N_i(x) \cap X_0 \subset A_\tau\}$ and $D_\tau = \bigcup_{i=0}^\infty \{x \in X_i \mid N_i(x) \cap B_\tau \neq \emptyset\}$. We have $C_\tau \cap D_\tau = \emptyset$ and $X \setminus (C_\tau \cup D_\tau) \subset \bigcup_{i=0}^\infty \{x \in X_i \mid N_i(x) \cap V_\tau^{X_0}(L) \neq \emptyset\}$.

Let us show that $\text{ex } C_\tau \supset \text{ex}_{X_0} A_\tau$. Choose any point $x \in \text{ex}_{X_0} A_\tau$. Then there exists $Z \subset X_0$ such that Z and $X_0 \setminus A_\tau$ are asymptotically disjoint and $x \in Z'$. Choose any $a > 0$. Since $k_i \rightarrow \infty$, there exists $n_0 \in \mathbb{N}$ such that $k_{n_0} \geq a$. Choose any $R > 0$ such that $d(Z \setminus B_R(x_0), (X_0 \setminus A_\tau) \setminus B_R(x_0)) \geq a + n_0$. Consider any points $z \in Z \setminus B_{R+n_0}(x_0)$ and $y \in \bigcup_{i=0}^\infty \{x \in X_i \mid N_i(x) \cap (X_0 \setminus A_\tau) \neq \emptyset\}$. We have that $d(z, y) \geq a$. Hence the sets Z and $\bigcup_{i=0}^\infty \{x \in X_i \mid N_i(x) \cap (X_0 \setminus A_\tau) \neq \emptyset\}$ are asymptotically disjoint and $x \in \text{ex } C_\tau$. Analogously we can show that $\text{ex } D_\tau \supset \text{ex}_{X_0} B_\tau$.

Now consider any $x \in C'_\tau \cap D'_\tau$. Then for every neighborhood V of x in the Higson compactification cX there exists two sequences (c_i) in $V \cap C_\tau$ and (d_i) in $V \cap D_\tau$ such that $0 < d(c_i, d_i) \leq r$ for some $r > 0$ and $c_i, d_i \in X \setminus B_i(x_0)$. Choose $n_0 \in \mathbb{N}$ such that $k_{n_0+1} > r$. So, $c_i, d_i \in \bigcup_{k=0}^{n_0} X_k$ and $\emptyset \neq \text{cl } V \cap (\bigcup_{k=0}^{n_0} X_k)' = \text{cl } V \cap X'_0$. Hence we have $x \in X'_0$. Moreover, $x \in X'_0 \setminus (\text{ex}_{X_0} A_\tau \cup \text{ex}_{X_0} B_\tau) = V_\tau^{X_0}(L)'$.

Denote $S_k(M) = \bigcup_{i=0}^\infty \{x \in X_i \mid N_{k,i}(x) \cap M \neq \emptyset\}$ for any $M \subset X_0$ and $k \in \mathbb{N}$. The set $S_1(M)$ we denote simply by $S(M)$. We have that $\nu X \setminus (\text{ex } D_\tau \cup \text{ex } C_\tau) = (X \setminus D_\tau)' \cap (X \setminus C_\tau)' \subset (X \setminus (D_\tau \cup C_\tau))' \cup (D'_\tau \cap C'_\tau) \subset (S(V_\tau^{X_0}(L)))'$.

Put $K_j = S(B_j(x_0))$ for each $j \in \mathbb{N}$. We have that the sets K_j and X_0 are asymptotically disjoint for each $j \in \mathbb{N}$. There exists $\sigma_j \in \omega^\omega$ such that $V_{\sigma_j}(X_0)' \cap K'_j = \emptyset$. Define $\sigma_0 \in \omega^\omega$ as follows $\sigma_0(0) = 0$ and $\sigma_0(i) = \max\{\sigma_j(i) \mid j \leq i\}$. We have that $\sigma_j \leq^* \sigma_0$ for each $j \in \mathbb{N}$. Hence $\text{cl } \bigcup_{j=1}^\infty K'_j \subset \nu X \setminus V_{\sigma_0}(X_0)'$.

Let us show that $\bigcap_{\tau \in \omega^\omega} (S(V_\tau^{X_0}(L)))' \cap V_{\sigma_0}(X_0)' \subset (S_2(L))'$. Choose any $x \notin (S_2(L))' \cup (\nu X \setminus V_{\sigma_0}(X_0)')$. Then there exists a subset $Z \subset X$ such that $x \in Z'$

and the sets $Z, S_2(L) \cup (X \setminus V_{\sigma_0}(X_0))$ are asymptotically disjoint. Let us consider the set $O(Z) \subset X_0$ defined as follows $O(Z) = \{y \in X_0 \mid \text{there exists } i \in \mathbb{N} \cup \{0\} \text{ and } x \in X_i \cap Z \text{ such that } d(y, x) \leq i\}$.

Suppose that there exist $r \in \mathbb{N}$ and two sequences (y_i) in $O(Z)$ and (l_i) in L such that $d(y_i, l_i) \leq r$ and $l_i, y_i \in X_0 \setminus B_i(x_0)$. For each i choose $x_i^{j_i} \in X_{j_i} \cap Z$ such that $d(x_i^{j_i}, y_i) \leq j_i$. Consider two cases:

1. there exists $n_0 \in \mathbb{N}$ such that $j_i \leq n_0$ for each $i \in \mathbb{N}$. Then $d(x_i^{j_i}, l_i) \leq n_0 + r$ and we obtain the contradiction with asymptotic disjointness of the sets Z and $S_2(L)$.

2. in the contrary case we can suppose that $j_i \geq r$ for each $i \in \mathbb{N}$ and $j_i \rightarrow \infty$. Then we have $d(x_i^{j_i}, l_i) \leq j_i + r \leq 2j_i$. Since $x_i^{j_i} \in X_{j_i}$, we have $x_i^{j_i} \in S_2(L) \cap Z$ and we obtain the contradiction again. So, the sets $O(Z)$ and L are asymptotically disjoint.

We can choose some $\tau \in \omega^\omega$ such that $O(Z)$ and $V_\tau^{X_0}(L)$ are asymptotically disjoint. Let us show that Z and $S(V_\tau^{X_0}(L))$ are asymptotically disjoint. Suppose the contrary. Then there exist $r \in \mathbb{N}$ and two sequences (z_i) in Z and (s_i) in $S(V_\tau^{X_0}(L))$ such that $d(z_i, s_i) \leq r$ and $z_i, s_i \in X_0 \setminus B_i(x_0)$. Consider two cases:

1. there exists $n_0 \in \mathbb{N}$ such that $z_i, s_i \in \cup_{i=0}^{n_0} X_i$. We can choose $y_i \in O(Z)$, $l_i \in V_\tau^{X_0}(L)$ for each $i \in \mathbb{N}$ such that $d(y_i, z_i) \leq n_0$ and $d(l_i, s_i) \leq n_0$. Then we have $d(y_i, l_i) \leq 2n_0 + r$ and we obtain the contradiction with the asymptotic disjointness of the sets $O(Z)$ and $V_\tau^{X_0}(L)$.

2. we can suppose that $z_i, s_i \notin \cup_{j=0}^{n_0} X_j$ where $k_n > r$ for each $n \geq n_0$. Then we have $z_i = s_i$ for each $i \in \mathbb{N}$ and we can choose $y_i \in O(Z) \cap V_\tau^{X_0}(L)$. Moreover, since Z and $X \setminus V_{\sigma_0}(X_0) \supset \cup_{j=1}^\infty K_j$ are asymptotically disjoint, we can assume that $d(y_i, x_0) \rightarrow \infty$. We obtain the contradiction again and the sets Z and $S(V_\tau^{X_0}(L))$ are asymptotically disjoint. Hence $x \notin (S(V_\tau^{X_0}(L)))'$ and we have $\cap_{\tau \in \omega^\omega} (S(V_\tau^{X_0}(L)))' \cap V_{\sigma_0}(X_0)' \subset (S_2(L))'$.

Put $V_A = \cup_{\tau \in \omega^\omega} \text{ex } C_\tau$ and $V_B = \cup_{\tau \in \omega^\omega} \text{ex } D_\tau$. Choose any point $x \in V_A \cap V_B$. Then there exist $\tau_1, \tau_2 \in \omega^\omega$ such that $x \in \text{ex } C_{\tau_1} \cap \text{ex } D_{\tau_2}$. Put $\tau = \max\{\tau_1, \tau_2\}$. Since $\text{ex } C_\tau \cap \text{ex } D_\tau = \emptyset$, we have that $x \in (\text{ex } C_{\tau_1} \setminus \text{ex } C_\tau) \cup (\text{ex } D_{\tau_2} \setminus \text{ex } D_\tau)$. Consider the case when $x \in (\text{ex } C_{\tau_1} \setminus \text{ex } C_\tau)$. Choose $n \in \mathbb{N}$ such that $A_{\tau_1} \setminus A_\tau \subset B_n(x_0)$. Then $x \in S(B_n(x_0))'$. The same we have in the case when $x \in (\text{ex } D_{\tau_2} \setminus \text{ex } D_\tau)$. We obtain that $V_A \cap V_B \subset \cup_{n \in \mathbb{N}} K'_n \subset \cup_{n \in \mathbb{N}} X_n \setminus V_n(X_0)'$.

We have that $V_A \supset O_A \supset A' \cap X'_0$ and $V_B \supset O_B \supset B' \cap X'_0$. Then there exists $\tau \in \omega^\omega$ such that $V_A \supset V_\tau(X_0)' \cap A'$ and $V_B \supset V_\tau(X_0)' \cap B'$. Moreover, we can suppose that $(S_2(L) \cap V_\tau(X_0))' \cap (A' \cup B') = \emptyset$ and $\sigma_0 \leq^* \tau$. Then $U_A^1 \cap U_B^2 = \emptyset$ where $U_A^1 = V_A \cap V_\tau(X_0)'$, $U_B^1 = V_B \cap V_\tau(X_0)'$. Then U_A^1 and U_B^1 are two disjoint open subset of νX such that $U_A^1 \supset V_\tau(X_0)' \cap A'$ and $U_B^1 \supset V_\tau(X_0)' \cap B'$. Since $\text{Ind}(\nu X \setminus V_\tau(X_0)') \leq 0$, there exists two open in $\nu X \setminus V_\tau(X_0)'$ disjoint sets U_A^2 and U_B^2 such that $U_A^2 \supset (\nu X \setminus V_\tau(X_0)') \cap A'$, $U_B^2 \supset (\nu X \setminus V_\tau(X_0)') \cap B'$ and $\nu X \setminus V_\tau(X_0)' = U_A^2 \cup U_B^2$. Since $V_\tau(X_0)'$ is clopen, U_A^2 and U_B^2 are open in νX .

The sets $U_A = U_A^1 \cup U_A^2$ and $U_B = U_B^1 \cup U_B^2$ are open disjoint subsets of νX such that $A' \subset U_A$, $B' \subset U_B$ and $\nu X \setminus (U_A \cup U_B) \subset (S_2(L) \cap V_\tau(X_0))'$.

Then $S_2(L) \cap V_\tau(X_0)$ is an asymptotic separator in X between A and B . Since L is a kernel of $S_2(L) \cap V_\tau(X_0)$, we have $\text{asInd}(S_2(L) \cap V_\tau(X_0)) < n$ by the inductive assumption and the theorem is proved.

3. In this section we prove the subspace and addition theorems.

Theorem 2. *Let X be a proper metric space and $Y \subset X$. Then $\text{asInd } Y \leq \text{asInd } X$.*

Proof. We shall apply induction with respect to $\text{asInd } X$. The case when $\text{asInd } X = -1$ is trivial. The case when $\text{asInd } X = 0$ follows from the equivalence $\text{asInd } X = 0$ and $\text{Ind } \nu X = 0$ for each metric proper space X [2].

Let us assume that we have proved the theorem for each X with $\text{asInd } X < n \geq 1$. Consider a proper metric space X with $\text{asInd } X = n$. Suppose that A and B are asymptotically disjoint subsets of Y and C is an asymptotic separator in X between V_A and V_B with $\text{asInd } C < n$ where V_A and V_B are asymptotic neighborhoods of A and B respectively such that V_A and V_B are asymptotically disjoint. We will build an asymptotic separator L in Y between A and B such that $\text{asInd } L < n$ using a construction from [4, Lemma 5.4].

Denote $D_0 = C$. Put $Z = Y \setminus (V_A \cup V_B)$. Let $D_k = (N_k(D_0) \cap Z) \setminus B_k(\cup_{i=0}^{k-1} D_i)$ for $k \in \mathbb{N}$. There exists a subset L_k of D_k such that L_k is k -discrete and for each $x \in D_k$ there exists $y \in L_k$ such that $d(x, y) \leq k$. Put $L = \cup_{i=1}^{\infty} L_i$. It was shown in [4] that L is an asymptotic separator in Y between A and B . It is easy to see that C is a kernel in $C \cup L$. Hence $\text{asInd}(C \cup L) < n$ and we have $\text{asInd } L < n$ by the inductive assumption. The theorem is proved.

Lemma 4. *Let X be a proper metric space and $f : cX \rightarrow [0, 1]$ is a continuous function. Consider any $a, b \in [0, 1]$ such that $a < b$. Then $(f^{-1}([a, b]) \cap X)' = (f^{-1}([0, b]) \cap X)' \cap (f^{-1}([a, 1]) \cap X)'$.*

Proof. Choose any $x \in (f^{-1}([0, b]) \cap X)' \cap (f^{-1}([a, 1]) \cap X)'$ and any neighborhood V of x in cX . Then there exist two sequences $(a_i), (b_i)$ in $V \cap X$ such that $f(a_i) \geq a$, $f(b_i) \leq b$, $d(a_i, b_i) \leq r$ for some $r > 0$ and $a_i, b_i \in X \setminus B_i(x_0)$ for each $i \in \mathbb{N}$. We can suppose that $f(a_i) \rightarrow c_1$ and $f(b_i) \rightarrow c_2$. Since $f|_X$ is slowly oscillating, then $c_1 = c_2 = c \in [a, b]$. Then we have $a < c \leq b$ or $a \leq c < b$. Consider the case $a \leq c < b$. There exists $n_0 \in \mathbb{N}$ such that $f(b_n) \in [a, b]$ for each $n \geq n_0$. Hence $\text{cl } V \cap (f^{-1}([a, b]) \cap X)' \neq \emptyset$ and $x \in (f^{-1}([a, b]) \cap X)'$. The proof is analogous in the case $a < c \leq b$. The inclusion $(f^{-1}([a, b]) \cap X)' \subset (f^{-1}([0, b]) \cap X)' \cap (f^{-1}([a, 1]) \cap X)'$ is trivial and the lemma is proved.

Theorem 3. *Let X be a proper metric space and $X = Y \cup Z$ where Y and Z are unbounded sets. Then $\text{asInd } X \leq \text{asInd } Y + \text{asInd } Z$.*

Proof. We shall apply induction with respect to $\text{asInd } Y + \text{asInd } Z$. If $\text{asInd } Y + \text{asInd } Z = 0$, then $\text{asInd } Y = \text{asInd } Z = 0$. We have that $\text{Ind } Y' = \text{Ind } Z' = 0$ [2]. Since $\nu X = Y' \cup Z'$, we have $\text{Ind } \nu X = 0$ [3, Theorem 2.2.7]. Then we have $\text{asInd } X = 0$ [2].

Assume that the theorem is proved for each Y and Z with $\text{asInd } Y \leq m \geq 0$, $\text{asInd } Z \leq l \geq 0$ and $m + l < n \geq 1$. Consider the case when $m + l = n$. Let A and B be asymptotically disjoint subsets of X . Then $A' \cap B' = \emptyset$. Choose a continuous function $f : \nu X \rightarrow [0, 1]$ such that $f(A') \subset \{0\}$ and $f(B') \subset \{1\}$ and extend it to a continuous function $g : cX \rightarrow [0, 1]$. Then the sets $g^{-1}[0, \frac{1}{3}] \cap X$ and $g^{-1}[\frac{2}{3}, 1] \cap X$ are asymptotically disjoint. We can choose an asymptotic separator L_1 in Y between $g^{-1}[0, \frac{1}{3}] \cap Y$ and $g^{-1}[\frac{2}{3}, 1] \cap Y$ such that $\text{asInd } L_1 < m$. Put $L_2 = g^{-1}[\frac{1}{3}, \frac{2}{3}] \cap Z$. Then $\text{asInd } L_2 \leq \text{asInd } Z \leq l$. We have that $\text{asInd } L_1 \cup L_2 \leq m - 1 + l < n$ by the inductive assumption.

Let us show that $L_1 \cup L_2$ is an asymptotic separator between A and B . It is easy to see that $K = g^{-1}[\frac{1}{3}, \frac{2}{3}] \cap X$ is an asymptotic separator between A and B in X . Put $V_1 = \text{ex}(g^{-1}[0, \frac{1}{3}] \cap X)$ and $V_2 = \text{ex}(g^{-1}(\frac{2}{3}, 1] \cap X)$. We have that V_1 and V_2 are disjoint open sets in νX such that $A' \subset V_1$ and $B' \subset V_2$. It follows from Lemma 4 that $\nu X \setminus K' \subset V_1 \cup V_2$. Since L_1 is an asymptotic separator in Y between

$g^{-1}[0, \frac{1}{3}] \cap Y$ and $g^{-1}[\frac{2}{3}, 1] \cap Y$, there exist two open in Y' disjoint sets O_1 and O_2 such that $O_1 \supset (g^{-1}[0, \frac{1}{3}] \cap Y)'$, $O_2 \supset (g^{-1}[\frac{2}{3}, 1] \cap Y)'$ and $Y' \setminus L'_1 = O_1 \cup O_2$. Put $V_2 = O_1 \setminus Z'$ and $U_2 = O_2 \setminus Z'$. Then the sets $V = V_1 \cup V_2$ and $U = U_1 \cup U_2$ are open disjoint subsets of νX such that $A' \subset V$, $B' \subset U$ and $\nu X \setminus (L_1 \cup L_2)' \subset V \cup U$. The theorem is proved.

4. We investigate the transfinite extension of asInd in this section. Let us recall the definition of the transfinite large inductive dimension trInd [3]: $\text{trInd } X = -1$ iff $X = \emptyset$; $\text{trInd } X \leq \alpha$ for an ordinal number if for every two disjoint closed subsets $A_1, A_2 \subset X$ there is a partition C with $\text{Ind } C \leq \alpha$.

Define the transfinite extension $\text{trasInd } X$ analogously: $\text{trasInd } X = -1$ if and only if X is bounded; $\text{trasInd } X \leq \alpha$ where α is an ordinal number if for every two asymptotically disjoint sets $A, B \subset X$ there is an asymptotic separator C with $\text{trasInd } C \leq \beta$ for some $\beta < \alpha$. Naturally we say $\text{trasInd } X = \alpha$ if $\text{trasInd } X \leq \alpha$ and it is not true that $\text{trasInd } X \leq \beta$ for some $\beta < \alpha$. We set $\text{trasInd } X = \infty$ if for each ordinal number α it is not true that $\text{trasInd } X \leq \alpha$.

The proof of the following theorem is the same as the one of Theorem 2.

Theorem 4. *Let X be a proper metric space and $Y \subset X$. Then $\text{trasInd } Y \leq \text{trasInd } X$.*

Let X be a metric proper space and $\{A_i | i \in \mathbb{N}\}$ is a countable family of subsets. We say that the family $\{A_i | i \in \mathbb{N}\}$ is *asymptotically discrete* if for each $i \in \mathbb{N}$ the sets A_i and $\cup_{j \neq i} A_j$ are asymptotically disjoint. We say that a proper metric space X is *asymptotically S -like* if X can be represented as the union of a sequence $X_1, X_2, \dots$ of subsets such that $\text{asInd } X_i \geq i$ and the family $\{X_i | i \in \mathbb{N}\}$ is asymptotically discrete. The class of all asymptotically S -like spaces we denote by $\mathcal{S}$.

Lemma 5. *If a proper metric space X is asymptotically S -like, then $\text{trasInd } X = \infty$.*

Proof. Suppose the contrary. Then there exists $X \in \mathcal{S}$ such that $\text{trasInd } X < \infty$. Put $\xi = \min\{\alpha \mid \text{there exists } X_\alpha \in \mathcal{S} \text{ such that } \text{trasInd } X_\alpha = \alpha\}$. Clearly, $\xi \geq \omega$ where ω is the first infinite ordinal number.

Choose any $X \in \mathcal{S}$ such that $\text{trasInd } X = \xi$. Let us represent X as the union $\cup_{i \in \mathbb{N}} X_i$ of subsets X_i such that $\text{asInd } X_i \geq i$ and the family $\{X_i | i \in \mathbb{N}\}$ is asymptotically discrete.

totically discrete. For each $i \in \mathbb{N}$ choose two asymptotically disjoint subsets A_i and B_i of X_i such that for each asymptotic separator L_i in X_i between A_i and B_i we have $\text{trasInd } L_i \geq i - 1$.

We build by induction sets $C_i \subset A_i$ and $D_i \subset B_i$ such that $d(C_i, \cup_{j=1}^i D_j) \geq i$ and $d(D_i, \cup_{j=1}^i C_j) \geq i$ for each $i \in \mathbb{N}$. Since A_1 and B_1 are asymptotically disjoint, there exists $r > 0$ such that $d(C_1, D_1) \geq 1$ where $C_1 = A_1 \setminus B_r(x_0)$ and $D_1 = B_1 \setminus B_r(x_0)$. Suppose we have built C_k and D_k for each $k \leq n \geq 1$. Build C_{n+1} and D_{n+1} . Since the sets $\cup_{j=1}^n X_j$ and X_{n+1} are asymptotically disjoint, there exists $r > 0$ such that $d((\cup_{j=1}^n X_j) \setminus B_r(x_0), X_{n+1} \setminus B_r(x_0)) \geq n + 1$. Since the sets A_{n+1} and B_{n+1} are asymptotically disjoint, there exists $t > 0$ such that $d(A_{n+1} \setminus B_r(x_0), B_{n+1} \setminus B_r(x_0)) \geq n + 1$. Put $s = \max\{r, t\}$ and $C_{n+1} = A_{n+1} \setminus B_s(x_0)$, $D_{n+1} = B_{n+1} \setminus B_s(x_0)$. It is easy to see that the sets $C = \cup_{i=1}^\infty C_i$, $D = \cup_{i=1}^\infty D_i$ are asymptotically disjoint and for each asymptotic separator L_i in X_i between C_i and D_i we have $\text{trasInd } L_i \geq i - 1$.

Choose any asymptotic separator L in X between C and D such that $\text{trasInd } L < \xi$. Since the family $\{X_i\}$ is asymptotically discrete, we have that $L_i = L \cap X_i$ is an asymptotic separator in X_i between C_i and D_i for each $i \in \mathbb{N}$. Then $\text{asInd } L_i \geq i - 1$. Thus, $L \in \mathcal{S}$ and we have a contradiction. The lemma is proved.

Lemma 6. *Let X be a proper metric space such that $\text{trasInd } X < \infty$. Then for each $x \in \nu X$ there exists a neighborhood V of x in cX such that $\text{asInd } V \cap X < \infty$.*

Proof. Suppose the contrary. Then there exists $x \in \nu X$ such that $\text{asInd } V \cap X = \infty$ for each neighborhood V of x in cX . Let us build by induction the sequence (L_i) of subsets of X and the sequence (V_i) of neighborhoods of x in cX such that the sets L_k and $V_k \cap X$ are asymptotically disjoint for each $k \in \mathbb{N}$, $L_{n+1} \subset V_n$, $\text{asInd } L_n = \infty$ and $V_1 \supset \dots \supset V_n \supset V_{n+1} \supset \dots$.

We have, particularly, that $\text{asInd } X = \infty$. There exist two asymptotically disjoint subsets $A, B \subset X$ such that $\text{asInd } L = \infty$ for each asymptotic separator L between A and B . We can assume that $x \notin B'$. Choose a continuous function $f : \nu X \rightarrow [0, 1]$ such that $f(x \cup A') \subset \{0\}$ and $f(B') \subset \{1\}$. Choose an asymptotic separator L_1 between the sets $f^{-1}[0, \frac{1}{3}] \cap X$ and $f^{-1}[\frac{2}{3}, 1] \cap X$ and put $V_1 = f^{-1}[0, \frac{1}{3}] \cap X$.

Assume we have built L_i and V_i for each $i \leq n \geq n + 1$. We have that $\text{trasInd } V_i$

∞ . So, there exist two asymptotically disjoint subsets C and D of V_n such that $\text{asInd } L = \infty$ for each asymptotic separator L between C and D . We can choose L_{n+1} and V_{n+1} as before. The sequences (L_i) and (V_i) are built. It is easy to check that the family L_i is asymptotically discrete. Put $L = \cup_{i=1}^{\infty} L_i$. So, $\text{trasInd } L = \infty$ by Lemma 5 and $\text{trasInd } X = \infty$ by Theorem 4.

Theorem 5. *Let X be a proper metric space such that $\text{trasInd } X < \infty$. Then $\text{asInd } X < \infty$.*

Proof. We can choose for each point $x \in \nu X$ a neighborhood V_x in cX such that $\text{asInd } V_x \cap X < \infty$. Since νX is compact, we have $\nu X \subset \cup_{i=1}^k V_i = V$ where each V_i is an open in cX set with $\text{asInd } V_i \cap X < \infty$. Then $\text{asInd } V \cap X < \infty$ by Theorem 3. Moreover, there exists $r > 0$ such that $X \subset V \cup B_r(x_0)$. Hence $\text{asInd } X < \infty$.

REFERENCES

1. M.Gromov, *Asymptotic invariants of infinite groups. Geometric group theory. v.2*, Cambridge University Press, 1993.
2. A.N. Dranishnikov, *On asymptotic inductive dimension*, JP J. Geom.Topol. **3** (2001), 239-247.
3. R.Engelking, *Dimension theory.Finite and infinite*, Heldermann Verlag, 1995.
4. A.Dranishnikov and M.M.Zarichnyi, *Universal spaces for asymptotic dimension (preprint)* (2001), 239-247.

DEPT. DE MATEMATICAS, FACULTAD DE CS. FISICAS Y MAT., UNIVERSIDAD DE CONCEPCION, CASILLA 160-C, CONCEPCION, CHILE
E-MAIL: TARASRADUL@YAHOO.CO.UK